

potential marker for monitoring patients with CHF and predicting their outcome.

Authors' affiliations

J George, S Patal, D Wexler, D Sheps, G Keren, A Roth, The Department of Cardiology, Tel Aviv Sourasky Medical Center, Tel Aviv, affiliated to the Tel Aviv University, Sackler School of Medicine, Tel Aviv Israel

Y Sharabi, E Peleg, Y Kamari, E Grossman, Hypertension Unit and Internal Medicine D, Sheba Medical Center, Tel Hashomer, Israel

REFERENCES

- Haluzik M, Parizkova J, Haluzik MM. Adiponectin and its role in the obesity-induced insulin resistance and related complications. *Physiol Res* 2004;**53**:123-9.
- Arita Y, Kihara S, Ouchi N, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 1999;**257**:79-83.
- Cnop M, Havel PJ, Utzschneider KM, et al. Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex. *Diabetologia* 2003;**46**:459-69.
- Weyer C, Funahashi T, Tanaka S, et al. Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 2001;**86**:1930-5.
- Hotta K, Funahashi T, Arita Y, et al. Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 2000;**20**:1595-9.
- Lindsay RS, Funahashi T, Hanson RL, et al. Adiponectin and development of type 2 diabetes in the Pima Indian population. *Lancet* 2002;**360**:57-8.
- Kumada M, Kihara S, Sumitsui S, et al. Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol* 2003;**23**:85-9.
- Ouchi N, Kihara S, Arita Y, et al. Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein, adiponectin. *Circulation* 1999;**100**:2473-6.
- Ouchi N, Kihara S, Funahashi T, et al. Reciprocal association of C-reactive protein with adiponectin in blood stream and adipose tissue. *Circulation* 2003;**107**:671-4.
- Matsubara M, Namioka K, Katayose S. Decreased plasma adiponectin concentrations in women with low-grade C-reactive protein elevation. *Eur J Endocrinol* 2003;**148**:657-62.
- Okamoto Y, Kihara S, Ouchi N, et al. Adiponectin reduces atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 2002;**106**:2767-70.
- Schrier RW, Abraham W. Hormones and hemodynamics in heart failure. *N Engl J Med* 1999;**341**:577-85.
- Anon. Heart Failure Society of America. HFSA guidelines for management of patients with heart failure caused by left ventricular systolic dysfunction: pharmacological approaches. *J Card Fail* 1999;**5**:357-82.
- Pugh PJ, Jones RD, Jones TH, et al. Heart failure as an inflammatory condition: potential role for androgens as immune modulators. *Eur J Heart Fail* 2002;**4**:673-80.
- Levine B, Kalman J, Mayer L, et al. Elevated circulating levels of tumor necrosis factor in severe chronic heart failure. *N Engl J Med* 1990;**323**:236-41.
- Lommi J, Pulkki K, Koskinen P, et al. Haemodynamic, neuroendocrine and metabolic correlates of circulating cytokine concentrations in congestive heart failure. *Eur Heart J* 1997;**18**:1620-5.
- Engeli S, Feldpausch M, Gorzelnik K, et al. Association between adiponectin and mediators of inflammation in obese women. *Diabetes* 2003;**52**:942-7.


IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.079129

Implantation of a biventricular defibrillator system in a patient with persistent left and absent right superior vena cava

A 61-year-old man with a previous myocardial infarction experienced heart failure symptoms (New York Heart Association functional class III). An echocardiogram revealed a severely impaired left ventricular ejection fraction (15%) under optimal medical treatment. Since he also showed a complete left bundle branch block and pronounced dyssynchrony on echocardiography he was referred for implantation of a biventricular implantable cardioverter-defibrillator (ICD) system.

After access to the left subclavian vein, a persistent left superior vena cava (LSVC) was encountered. Subsequently, a venogram also revealed the absence of a right superior vena cava (panel A: AV, anomalous vein). The persistent LSVC drained into the notably enlarged coronary sinus (CS, panel B). Therefore, all three leads had to be implanted through the coronary sinus. The proximal shock coil was located within the coronary sinus. The left ventricular electrode (LV) had to be implanted through the posterior cardiac vein and could be positioned at a favourable position in a lateral branch (panel C: A, atrial lead in the right atrial appendage; RV, right ventricular defibrillation lead; panel D: x ray with the successfully implanted cardiac resynchronisation therapy-defibrillator system). The patient could be successfully defibrillated with 12 J and experienced significant improvement in heart failure symptoms briefly after the initiation of biventricular pacing. The persistence of the left superior vena cava is the most commonly encountered congenital anomaly of the venous system with a prevalence of 0.3-1%. This



case illustrates the feasibility of successfully implanting a biventricular ICD system through a persistent left superior vena cava and the coronary sinus.

**C Sticherling
S Osswald
csticherling@uhbs.ch**